

Phase fluctuations and Non-Fermi Liquid Properties of 2D Fermi-system with attraction

Valery P. Gusynin¹, Vadim M. Loktev¹, Rachel M. Quick² and Sergei G. Sharapov^{2,†}

¹*Bogolyubov Institute for Theoretical Physics,
252143 Kiev, Ukraine*

²*Department of Physics, University of Pretoria,
0002 Pretoria, South Africa*

The effect of static fluctuations in the phase of the order parameter on the normal and superconducting properties of a 2D system with attractive four-fermion interaction has been studied. Analytic expressions for the fermion Green function, its spectral density and the density of states are derived. The resultant single-particle Green function clearly demonstrates non-Fermi liquid behavior. The results show that as the temperature increases through the 2D critical temperature the width of the quasiparticle peaks broadens significantly. At the same time one retains the gap in quasiparticle spectrum. The spectral density for the dynamical fluctuations can also be obtained. Clearly the dynamical fluctuations fill the gap giving the observed pseudogap behaviour.

[†] On leave from Bogolyubov Institute for Theoretical Physics of the National Academy of Sciences of Ukraine, 252143 Kiev, Ukraine

Invited paper presented at the Second International Conference on
New Theories, Discoveries, and Applications of Superconductors and Related Materials
May 31– June 4, 1999, Las Vegas, Nevada, USA

It is known [1] that the phase fluctuations are especially important in low dimensional superconductors with small carrier density (such as the high- T_c oxide superconductors). In particular there is some experimental evidence that the transition into superconducting state in high- T_c cuprates is driven by phase fluctuations.[2] The microscopic theory developing this scenario [1] has been proposed in [3].

The main quantity of interest in the present report is the one-fermion Green's function and the associated spectral function $A(\omega, \mathbf{k}) = -(1/\pi)\text{Im}G(\omega + i0, \mathbf{k})$. The second quantity, being proportional to the intensity of the angle-resolved photoemission spectrum (ARPES) [4], encodes information about the quasiparticles. Following the approach of Ref.[3] the Green's function for the charged (physical) fermions, $G(\omega, \mathbf{k})$ is given by the convolution (in momentum space) of the propagator for neutral fermions which has a gap $\rho \neq 0$ and the Fourier transform of the phase correlator $\langle \exp(i\theta(x)/2) \exp(-i\theta(0)/2) \rangle$.

We demonstrate that the quasiparticle spectral function broadens considerably when passing from the superconducting to the normal state as observed experimentally [4]. More importantly the phase fluctuations result in non-Fermi liquid behavior of the system both below and above T_{BKT} , where T_{BKT} is the temperature of Berezinskii-Kosterlitz-Thouless (BKT) superconducting transition in 2D.

In the frequency-momentum representation the Green function of charged fermions can be written (see details in [5]) as

$$G(i\omega_n, \mathbf{k}) = T \sum_{m=-\infty}^{\infty} \int \frac{d^2p}{(2\pi)^2} \sum_{\alpha, \beta=\pm} P_{\alpha} \mathcal{G}(i\omega_m, \mathbf{p}) P_{\beta} D_{\alpha\beta}(i\omega_n - i\omega_m, \mathbf{k} - \mathbf{p}), \quad (1)$$

where $\mathcal{G}(i\omega_m, \mathbf{k})$ is the Green function of neutral fermions; $P_{\pm} \equiv (1/2)(\hat{I} \pm \tau_3)$ are the projectors; $\hat{I}$, τ_3 are unit and Pauli matrices; $D_{\alpha\beta}(i\Omega_n, \mathbf{q})$ is the Fourier transform of the correlator of the phase fluctuations $\langle \exp[i\alpha\theta(x)/2] \exp[-i\beta\theta(0)/2] \rangle$ ($x = \tau, \mathbf{r}$ with the imaginary time τ and 2D space coordinate $\mathbf{r}$); $\omega_n = (2n+1)\pi T$, $\Omega_n = 2\pi nT$ are fermion (odd) and boson (even) Matsubara frequencies.

One can show that $D_{+-} = D_{-+} = 0$ which retains the symmetric form of the propagator G is accordance to the Coleman-Mermin-Wagner-Hohenberg theorem and the properties of (1) are entirely determined by $D \equiv D_{++} = D_{--}$.

The general form of the phase correlator is

$$D(t, \mathbf{r}) = \exp(-\gamma t) \left(\frac{r}{r_0} \right)^{-\frac{T}{8\pi J}} \exp \left(-\frac{r}{\xi_+(T)} \right), \quad (2)$$

where γ is the phenomenological decay constant; $J(T)$ is the superfluid stiffness [3]; r_0 is the scale of algebraic decay of the correlations in the BKT phase ($T < T_{\text{BKT}}$); $\xi_+(T)$ is the correlation length for $T > T_{\text{BKT}}$ ($\xi_+ = \infty$ for $T \leq T_{\text{BKT}}$). Note that t is the real time, so that (2) is the retarded Green function. The power decay factor in Eq. (2) is related to the spin-wave phase fluctuations, while the exponent with ξ_+ is responsible for the vortex excitations, which are present only above T_{BKT} .

It is remarkable that the static case ($\Omega_n = \gamma = 0$) can be studied analytically and one obtains the retarded Green function [5]

$$G(\omega, \mathbf{k}) = -\frac{Cm\xi_+^{2\alpha}}{2\pi\alpha} \left[\frac{\mathcal{A}}{(v_1v_2)^\alpha} F_1 \left(\alpha, \alpha, \alpha; \alpha + 1; \frac{v_1 - 1}{v_1}, \frac{v_2 - 1}{v_2} \right) + (\sqrt{\omega^2 - \rho^2} \rightarrow -\sqrt{\omega^2 - \rho^2}) \right], \quad (3)$$

where F_1 is the Appell's function of two variables,

$$\begin{aligned} \alpha &= 1 - \frac{T}{16\pi J(T)}, \quad C = \frac{4\pi\Gamma(\alpha)}{\Gamma(1-\alpha)} \left(\frac{2}{r_0} \right)^{2\alpha-2}, \quad \mathcal{A} = \frac{1}{2} \left(\tau_3 + \frac{\omega}{\sqrt{\omega^2 - \rho^2}} \right), \\ v_{1,2} &= m\xi_+^2 (v_0 \pm \sqrt{\mathcal{D}}), \quad \mathcal{D} = v_0^2 + \frac{2}{m\xi_+^2} (\mu + \sqrt{\omega^2 - \rho^2}), \\ v_0 &= \frac{k^2\xi_+^2 + 1}{2m\xi_+^2} - \mu - \sqrt{\omega^2 - \rho^2}, \end{aligned} \quad (4)$$

m is the mass of fermions.

For $T < T_{\text{BKT}}$ near the quasiparticle peaks when $\omega \approx \pm \sqrt{(\mathbf{k}^2/2m - \mu)^2 + \rho^2}$ one can derive from (3) that

$$G(\omega, \mathbf{k}) \sim -\Gamma^2(\alpha) \left(\frac{2}{mr_0^2} \right)^{\alpha-1} \frac{\mathcal{A}}{[-(\mu + \sqrt{\omega^2 - \rho^2})]^\alpha} \frac{\Gamma(2\alpha - 1)}{\Gamma^2(\alpha)} \frac{1}{(1 - z_1)^{2\alpha-1}}, \quad (5)$$

where $z_1 \equiv (k^2/2m)/(\mu + \sqrt{\omega^2 - \rho^2}) \simeq 1$. It is seen that Eq. (5) is evidently nonstandard since it contains a branch cut, not pole. The latter in its turn corresponds to the non-Fermi liquid behavior of the system as whole. It must be underlined that non-Fermi liquid peculiarities are tightly related to the charge (i.e. observable) fermions only – the Green function of neutral ones has typical (pole type) BCS form. Besides, because the parameter α is a function of T (see (4)) the non-Fermi liquid behavior is becoming more pronounced as temperature increases and is preserved until ρ vanishes.

The spectral function $A(\omega, \mathbf{k})$ associated with $G(\omega, \mathbf{k})$ has been studied in [5]. Here we sketch the major features. *i)* The quasiparticle peaks of the spectral density have a finite temperature dependent width which decreases with decreasing T in the superconducting state. The sharpening of the peaks with decreasing T is seen experimentally [4] and this is in a striking difference from the BCS temperature independent “pile-up”. *ii)* For $T > T_{\text{BKT}}$ the width of peaks broadens dramatically which is also observed experimentally [4]. *iii)* In the static approximation ($\Omega_n = \gamma = 0$) we obtained two extra peaks which are present for $T \neq 0$. The gap in the excitation spectrum remains nonfilled even above T_{BKT} . *iv)* The spectral density for the dynamical fluctuations, i.e. when $\Omega_n \neq 0$ modes are taken into account can also be obtained. The extra peaks seen in the static case disappear. Clearly

the dynamical fluctuations fill the gap giving the observed pseudogap behaviour [4] where the strength of the filling depends on γ .

Acknowledgements

R.M.Q and S.G.Sh acknowledge the financial support of the NRF, Pretoria.

References

- [1] V. Emery, S.A. Kivelson, *Nature* **374**, 434 (1995); *Phys. Rev. Lett.* **74**, 3253 (1995).
- [2] J. Corson, R. Mallozzi, et al., *Nature* **398**, 221 (1999).
- [3] V.P. Gusynin, V.M. Loktev, S.G. Sharapov, *JETP Lett.* **65**, 182 (1997); *JETP* **88**, 685 (1999); V.M. Loktev, S.G. Sharapov, *Cond. Mat. Physics (Lviv)* No.11, 131 (1997) (cond-mat/9706285); R.M. Quick, S.G. Sharapov, *Physica C* **301**, 262 (1998); V.M. Loktev, R.M. Quick, S.G. Sharapov, *Physica C* **314**, 233 (1999), cond-mat/9904126, to appear in *Low Temp. Phys.* (2000).
- [4] M. Randeria and J.C. Campuzano, in *Proceedings of the International School of Physics*, “Enrico Fermi”, Varenna, 1997 (North Holland, New York) (cond-mat/9709107).
- [5] V.P. Gusynin, V.M. Loktev, S.G. Sharapov, *JETP Lett.* **69**, 141 (1999); cond-mat/9811207, to appear in *JETP* (2000).